

To be published in *HST and the High Redshift Universe (37th Herstmonceux Conference)*, eds. N.R. Tanvir, A. Aragon-Salamanca, J.V. Wall, 1996.

THE ULTRAVIOLET MORPHOLOGY OF GALAXIES

ROBERT W. O'CONNELL, PAMELA MARCUM

*University of Virginia,
Charlottesville, VA 22903-0818, USA*

Optical band images of distant ($z \gtrsim 0.5$) galaxies, such as those of the Hubble Deep Field, record light from the rest-frame vacuum ultraviolet ($\lambda < 3000 \text{ \AA}$). Because the appearance of a galaxy is a very strong function of wavelength, and especially so in the UV, evolutionary studies of distant galaxies can be seriously influenced by a “morphological k-correction” effect. We use images obtained by the Ultraviolet Imaging Telescope during the *Astro* missions to explore the extent of this effect and intercompare far-UV with optical morphologies for various types of galaxies.

The morphology of a distant galaxy is often our first index to its evolutionary state. Unfortunately, there is a host of technical complications which afflict the apparent morphologies of distant galaxies. These can conspire to make the familiar look unfamiliar and vice-versa.

Two of these difficulties are well known: (i) reduced spatial resolution; and (ii) the rapid fading of surface brightness with redshift, $SB \sim (1+z)^{-n}$, where $n = 3-5$, depending on the method used to characterize surface brightness. These two effects are relatively simple to model, and their consequences for ground-based and HST imaging have recently received much attention^{1 2 3 4 5}. Here, we wish to focus on a third difficulty, which can be called the “morphological k correction”—i.e. the fact that the appearance of galaxies is a strong function of wavelength. The photons recorded on exposures of distant galaxies in optical bands, e.g. in the Hubble Deep Field, often originate in the vacuum ultraviolet in the restframe. It has been known for many years that the appearance of galaxies in the UV can be very different than at the optical wavelengths for which the classical morphological typing systems were established^{6 7}. In the UV, hot stars are emphasized and cool ones are suppressed. This provides an important opportunity to distinguish stellar populations, but it also means that morphological analyses must take into account a strong UV/optical k-correction.

It has been hard to quantify this effect because few vacuum-UV images of nearby galaxies were available before 1990. Since then, several hundred objects have been observed in the UV by HST, the FOCA balloon experiment⁸, and the Ultraviolet Imaging Telescope⁹ (UIT) on the two *Astro* missions. Maos et al.¹⁰ have recently published an atlas of 110 nearby galaxy nuclei observed at




Figure 1: Dust does not dominate the UV appearance of normal disk galaxies, even in edge-on systems like this one.

2300 Å in the 22'' field of the HST Faint Object Camera. We¹¹ are preparing a UV/optical atlas which compares the UV and optical morphologies of 27 nearby galaxies using UIT data mainly from the *Astro-1* mission. A later paper will add 45 galaxies observed during *Astro-2*. UIT's field of view is 40' in diameter and produced typical resolution of 3'' FWHM; the two principal (solar-blind) imaging bands were centered at 1500 Å ("far-UV") and 2500 Å ("mid-UV"). UIT's characteristics are well suited to the study of morphologies of nearby galaxies. Some of this data has already been used to simulate the combined effects of the k-correction, reduced resolution, and reduced surface brightness on observations of distant galaxies^{5 12}.

Here we show a few of the comparisons from the UIT Atlas and briefly mention the effects they illustrate. Data for a number of spirals shows that dust in normal disks does not strongly affect UV morphologies (Fig. 1). However, spirals do shift to later Hubble types in the UV (Fig. 2). The k-correction combined with reduced resolution and bias toward high SB regions can turn a single normal galaxy into an irregular object with "interacting" companions (Fig. 3). Disturbed dust layers can have drastic effects on UV light (e.g. M82, not shown); for instance, an interacting pair of galaxies can turn into an isolated galaxy (Fig. 4). Barred galaxies often become unbarred (e.g. M83⁶). The general trend found is from normal to abnormal. To put this another way, the incidence of abnormal galaxies in the restframe UV appears to be larger than it really is.

References

1. S.P. Driver, R.A. Windhorst, and R.E. Griffiths, *ApJ*, 453, 48, 1995.
2. S.J. Lilly, O. Le Fevre, D. Crampton, F. Hammer, and L. Tresse, *ApJ*, 455, 50, 1995.




Figure 2: Hubble types for spirals become later in the UV. In the far-UV (not shown), M81 becomes a nearly “empty” ring.




Figure 3: Effects of k-correction and reduced resolution, corresponding to observations at $z \sim 1$ with $0.4''$ FWHM. Bright associations have become “companions”.




Figure 4: Case of a disappearing perturber.

3. R.G. Abraham, N.R. Tanvir, B.X. Santiago, R.S. Ellis, K. Glazebrook, and S. van den Bergh, MNRAS, in press.
4. A. Dressler, this conference.
5. M. Giavalisco, J. Livio, R.C. Bohlin, F.D. Macchetto, and T.P. Stecher, AJ, 112, 369, 1996.
6. R.C. Bohlin, R.H. Cornett, J.K. Hill, A.M. Smith, and T.P. Stecher, ApJ, 274, L53, 1983.
7. C.R. King and R. S. Ellis ApJ, 288, 456, 1985.
8. A. Blecha, M. Golay, D. Huguenin, D. Reichen, and D. Bersier, A&A, 233, L9, 1990.
9. T. P. Stecher et al. ApJ, 395, L1, 1992.
10. D. Maoz, A.V. Filippenko, L.C. Ho, F.D. Macchetto, H.-W. Rix, and D.P. Schneider, ApJS, in press.
11. P. Marcum, R.W. O'Connell, W. Freedman, B. Madore, G. Bothun, K.P. Cheng, P.M. Hintzen, E.P. Smith, R.C. Bohlin, S.G. Neff, M.S. Roberts, A.M. Smith, and T.P. Stecher, in preparation for ApJS.
12. R.C. Bohlin, R.H. Cornett, J.K. Hill, R.S. Hill, W.B. Landsman, R.W. O'Connell, S.G. Neff, A.M. Smith, and T.P. Stecher, ApJ, 368, 12, 1991.